

MISTI 68.35.47

UDC 633.317

https://doi.org/10.52269/22266070_2024_2_55

**THE EFFECT OF FOLIAR TREATMENT OF ALFALFA CROPS DURING
VEGETATION USING BIOPREPAREATIONS ON SEED YIELD**

Kalin A.K.* – PhD student, "8D08101–Agronomy" educational program, Sh.Ualikhanov Kokshetau University, Republic of Kazakhstan.

Sagalbekov U.M. – Doctor of Agricultural Sciences, Professor of the Department of agriculture and bioresources, Sh.Ualikhanov Kokshetau University, Republic of Kazakhstan.

Auzhanova M.A. – Candidate of Agricultural Sciences, acting Associate Professor of the Department of agriculture and bioresources, Sh.Ualikhanov Kokshetau University, Republic of Kazakhstan.

Ualiyeva G.T. – PhD student, "8D08101 – Agronomy" educational program, Kokshetau experimental production farm, Akmola region, Shagalaly village, Republic of Kazakhstan.

The article presents the results of a study on the influence of biopreparations BioSleepBW+Foliar, OrgamicaS+Foliar, (OrganitP+OrganitN+Biodux+Foliar), and Foliar on the seed yield of alfalfa when applied as foliar treatments at the beginning of flowering. The research on alfalfa cultivation was conducted using five different experimental variants. The experiments were carried out on ordinary chernozem soil. The Hydrothermal Coefficient (HTC) values in 2022-2023 were 0,73 – 0,76, respectively, indicating dry conditions. During the budding and flowering periods of alfalfa plants, there was a precipitation deficit, which negatively affected seed yield.

The composition of the biopreparations includes a complex of biologically active polyunsaturated fatty acids of the fungus Mortierella alpina with a consumption rate of 3,0-10,0 ml/ha and spores of the strains Beauveria bassiana, Bacillus megaterium-amyloliquefaciens with consumption rates of 1-3 l/ha, with a working solution consumption of 300 l/ha.

The research established that the applied biopreparations improve the sowing quality of the seeds. The weight of 1000 seeds remains largely unchanged, serving as a stable indicator, while the agricultural validity of the planting material increases from 79.2% to 84%. This ultimately leads to a 10-20% increase in alfalfa seed yield.

The most effective biopreparations influencing seed productivity are Orgamica S+Foliar with a seed yield of 2,0 c/ha, Foliar with 1,86 c/ha, and the OrganitP+OrganitN+Biodux+Foliar complex with 1,82 c/ha, which exceed the control variant by 0,37, 0,28, and 0,05 c/ha, respectively.

Key words: alfalfa; biopreparations; foliar treatment; economic validity; seed yield.

**ЖОҢЫШҚА ӨСІНДІЛЕРІН БИОПРЕПАРТАРМЕН ВЕГЕТАЦИЯ БОЙЫНША
ТАМЫРДАН ТЫС ӨНДЕУДІҢ ТҮҚЫМ ӨНІМДІЛІГІНЕ ӘСЕРІ**

Калин А.К.* – "8D08101 – Агрономия" мамандығы бойынша докторантуралың білім алушысы, Ш. Уәлиханов атындағы Көкшетау университеті, Көкшетау қ., Қазақстан Республикасы.

Сагалбеков У.М. – ауыл шаруашылығы ғылымдарының докторы, ауыл шаруашылығы және биоресурстар қафедрасының профессоры, Ш. Уәлиханов атындағы Көкшетау университеті, Көкшетау қ., Қазақстан Республикасы.

Аужанова М.А. – ауыл шаруашылығы ғылымдарының кандидаты, ауыл шаруашылығы және биоресурстар қафедрасының қауымдастырылған профессорының м. а., Ш. Уәлиханов атындағы Көкшетау университеті, Көкшетау қ., Қазақстан Республикасы.

Үәлиева Г.Т. – "8D08101-Агрономия" мамандығының докторантты, Көкшетау тәжірибелік-өндірістік шаруашылығы, Ақмола облысы Шагалалы ауылы, Қазақстан Республикасы.

Бұл мақалада Biosleepbw+Foliar, OrgamicaS+Foliar, (OrganitP+OrganitN+Biodux+Foliar) және Foliar биопрепараттарының вегетация бойынша тамырдан тыс өнделген жоңышқа түқымдарының өнімділігіне әсері бойынша зерттеу нәтижелері қарастырылады. Жоңышқа өсіру бойынша зерттеулер тәжірибелің бес нұсқасы бойынша қаралайым қара топырақта жүреілді. 2022-2023 жылдардағы гидротермиялық коэффициенттің (ГТК) шамалары тиісінше 0,73–0,76 құрады, бұл құргақ болып табылады. Осы уақытта бүршіктену және ғулдену кезеңінде жауын-шашынның жетіспеушілігі байқалды, бұл түқым өнімділігіне теріс етті.

Биопрепараттардың құрамы Mortierella alpina санырауқұлақтарының биологиялық белсенеді полиқанықпаган май қышқылдары кешенінен құрастырылған, тұтыну нормасы 3,0 – 10,0 мл/га және Beauveria bassiana, Bacillus megaterium – amyloliquefaciens штаммдарының спораларымен, шығын нормалары бойынша 1-3 л/га, жұмыс ерітіндісінің шығыны-300 л/га.

Зерттеулер көрсөткендегі, қолданылатын биопрепараттар тұқымның себу сапасын жақсартады. 1000 тұқымның массасы практикалық ең тұрақты белгі ретінде өзгермейді, сонымен бірге тұқымның шаруашылық жарамдылығы 79,2% – дан 84% –ға дейін артады, нәтижесінде жонышқа тұқымының өнімділігі 10-20% –ға артады. Тұқым өнімділігі бойынша ең тиімділері *Organica S+Foliar* (2,0 ү/га), *Foliar* (1,86 ү/га) препараттары және *OrganitP+OrganitN+Biodux+Foliar* (1,82 ү/га) болды. Өнімділік көрсөткіштері бақылау нұсқасынан сәйкесінше 0,37, 0,28 және 0,05 ү/га-ға артық.

Түйінді сөздер: жонышқа; биопрепараттар; жапырақты өңдеу; шаруашылық жарамдылығы; тұқым өнімділігі.

ВЛИЯНИЕ ВНЕКОРНЕВОЙ ОБРАБОТКИ ПОСЕВОВ ЛЮЦЕРНЫ ПО ВЕГЕТАЦИИ БИОПРЕПАРАТАМИ НА УРОЖАЙНОСТЬ СЕМЯН

Калин А.К.* – обучающийся докторанттуры по специальности «8D08101 – Агрономия», Кокшетауский университет имени Ш. Уалиханова, г. Кокшетау, Республика Казахстан.

Сагалбеков У.М. – доктор сельскохозяйственных наук, профессор кафедры «Сельское хозяйство и биоресурсы», Кокшетауский университет имени Ш. Уалиханова, г. Кокшетау, Республика Казахстан.

Аужанова М.А. – кандидат сельскохозяйственных наук, и.о. ассоциированного профессора кафедры сельское хозяйство и биоресурсы, Кокшетауский университет имени Ш. Уалиханова, г. Кокшетау, Республика Казахстан.

Уалиева Г.Т. – докторант специальности «8D08101 – Агрономия», Кокшетауское опытно-производственное хозяйство, Акмолинская область с. Шагалапы, Республика Казахстан.

В статье приведены результаты исследования по влиянию биопрепаратов *BioSleepBW+Foliar*, *OrganicaS+Foliar*, (*OrganitP+OrganitN+Biodux+Foliar*) и *Foliar* на урожайность семян люцерны при внекорневой обработке посевов в период начала цветения. Исследования по возделыванию люцерны проводились по пяти вариантам опыта. Опыты проводились на черноземе обыкновенном. Величины гидротермического коэффициента (ГТК) в 2022-2023 годы составляли 0,73–0,76, соответственно, что являются засушливыми. В период бутонизации и цветения растений люцерны наблюдался недобор осадков, что отрицательно сказалось на урожайность семян.

Состав биопрепаратов представляют собой комплекс биологически активных полиненасыщенных жирных кислот гриба *Mortierella alpine* с нормой расхода 3,0-10,0 мл/га и спорами штаммов *Beauveria bassiana*, *Bacillus megaterium*- *amyloliguefaciens* с нормами расходов 1-3 л/га, расход рабочего раствора – 300 л/га.

Исследованиями установлено, что применяемые биопрепараты улучшают посевные качества семян. Масса 1000 семян практически не меняется как наиболее стабильный признак, и при этом хозяйственная годность посевного материала повышается с 79,2% до 84%, что в конечном итоге приводит к увеличению урожайности семян люцерны на 10-20%. Наиболее эффективными биопрепаратами по влиянию на продуктивность семян являются *Organica S+Foliar* с урожайностью семян 2,0 ү/га, *Foliar*-1,86 ү/га и комплекс *OrganitP+OrganitN+Biodux+Foliar*-1,82 ү/га, что превышает контрольный вариант на 0,37, 0,28 и 0,05 ү/га соответственно.

Ключевые слова: люцерна; биопрепараты; внекорневая обработка; хозяйственная годность; урожайность семян.

Introduction. The development of animal husbandry for 2020-2030 directly depends on feed production. The priority task of the industry today is to create a cost-effective highly productive base with high-quality dry matter feed, which should contain 15-25% protein and 8-11 MJ of metabolic energy. All these measures can be carried out with the help of state support for the implementation of long-term programs for the development of forage production, in particular, the development of seed production, raising the yield of forage crops, the acquisition of modern forage harvesting and forage preparation equipment [1, p.44].

The irrational structure of the acreage with 5-8% of perennial grasses, and 70-80% of cereals led to a decrease in the production of bulky and green feeds. As a result, it led to a decrease in 1 kg of dry matter of feed units less than 0,40 protein content – 8-9% and metabolic energy 7-8 MJ in the produced feeds [2, p.50;3, p.17].

Intensification of animal husbandry is impossible without expanding the acreage occupied by perennial grasses. It reduces the cost of producing grain feeds, and solves the problems of producing high-protein feeds with a high content of carotene, various vitamins, mineral salts, and trace elements [4, p.2705;5, p.380].

To improve the livestock feed base, the areas occupied for field feed production should account for 20-30% of the total sown area of the republic. Currently, Kazakhstan has 5,0 million hectares of hayfields,

186 million hectares of pastures, and 2,5 million hectares of arable land occupied for fodder crops, which is 12% of the total sown area [6, p.139;7, p.400].

The restoration and development of feed production directly depend on the cultivation of perennial grasses, which will improve the structure of acreage, and reduce the industry's financial, technical, and energy costs by 25-35%. To this end, it is necessary to create highly productive varieties of perennial grasses, with the help of which it is possible to solve the problems of providing livestock with high-grade feed. For each region with its natural and climatic conditions, the choice of specific perennial grass varieties is crucial for obtaining sustainable harvests [8, p.4].

Animal husbandry needs a solid feed base that provides high-protein feeds. Solving this problem by sowing perennial grasses, especially alfalfa, is possible. Alfalfa is of great forage importance among perennial forage grasses, due to its high content of protein, carotene, various vitamins, mineral salts, and trace elements. Its crops can be used as pastures for canned feed (silage, haylage), hay harvesting, and grass flour. In addition, compared with other legumes used in forage production, alfalfa has higher winter hardiness, resistance to weed infestation, and the ability to grow a large green mass in one season [9, p.110].

Alfalfa is an easily digestible perennial plant for animals with a high content of protein, vitamins, trace elements, and all kinds of acids. The plant saturates the body with the necessary amount of iron, fluorine, potassium, magnesium, and calcium [10, p.60].

Alfalfa is a biological meliorant, which is an excellent precursor for all crops. Alfalfa frees the soil from harmful microorganisms and enriches it with useful ones [11,p.128].

Alfalfa is used for green fodder, grass flour, hay, and haylage. During the flowering period, 20-22% of crude protein is concentrated in its aboveground mass. The digestibility of alfalfa protein is 77% versus 65-75% for other perennial grasses. Drought resistance, deep root system, symbiotic nitrogen fixation, rapid regrowth, all these signs are characteristic of alfalfa changeable [12, p.110].

Alfalfa is provided with micro- and macroelements due to the fixation of atmospheric nitrogen with the support of nodule bacteria. But in natural conditions, alfalfa uses only 10-35% of its nitrogen-fixing capacity, which leads to a lack of nitrogen in the soil [13, p.182].

The use of pesticides in the cultivation of legumes negatively affects plant growth and soil fertility. Improvement of indicators of physical properties of soils, and leaf-stem mass of plants is possible with the introduction of bacterial fertilizers and seed treatment before sowing with elements of organic farming [14, p.50].

Nitrogen is a component of many legumes that is essential for plant growth. A significant part of nitrogen is contained in the atmosphere and its use is important to prevent nitrogen starvation. However, the ability to fix atmospheric nitrogen by nodule bacteria is limited. In this case, the application of biologized preparations based on *Rhizobium* strains increases the content of total organic carbon, nitrogen, phosphorus, and potassium in the soil, increases plant biomass, and increases the physical properties of soils [15, p.56;16, p.37;17, p.67].

The growth of the world population and the need for high-calorie quality products require the search for new ways to solve the food supply of the population with agricultural products obtained based on an ecological, biologized component of the crop industry [18, p.1062].

There is a need to practice the transition from traditional farming to organic farming. According to the data of the International Federation of Ecological Agricultural Movement (IFOAM) and the Research Institute (FIBL) for 2023, the land areas in the world allocated for organic agriculture for this period are: Europe – 17,8 million hectares, Oceania – 36 million hectares, Africa – 2,7 million hectares, North America – 3,5 million hectares, Latin America America – 9,9 million hectares, Asia – 6,5 million hectares. The largest markets for organic products are in the USA – 48,7 billion euros, Germany – 16 billion euros, France – 12,8 billion euros [19, p.2].

There are potential opportunities for Kazakhstan to provide the domestic sector with high-quality feed and breakthrough areas in the export of products outside the Republic. The global feed producer (APEHF), where the European Union and the United States are the leaders, exporting 3 and 5 million tons of hay per year, offers Kazakhstan, in particular, large-scale cultivation of alfalfa for export in the form of granular feed and pressed hay to the countries of the Middle East and China, where a ton of dry alfalfa hay costs 200-250 thousand tenge. In this case, the parameter determining the price is the high-protein quality of alfalfa feed with a protein content of 15% and higher[20,p.10].

Improving the technology of growing alfalfa in rainfed will always remain a priority in feed production. In this regard, one of the effective measures to increase the seed productivity of alfalfa is the use of biological products that help increase the yield of green mass and seeds, which made it possible to formulate the goal of our research – to study the effect of biological products on the yield of alfalfa seeds in the conditions of the Akmola region. The main objective of the study is to study the effect of biological products with seed inoculation and during the growing season on the growth, development and yield of alfalfa seeds.

Materials and methods of research. Field experiments using biological products were conducted in the hill-plain zone in Kokshetau Experimental Production Farm LLP, Akmola region, Zerendinsky district,

Shagalaly village. In three-fold repetition. The sowing method is ordinary and wide-row, row spacing is 70 cm. The agricultural technology in the experiments is zonal. The area of the experimental plot is 20 m², the placement of plots is randomized. The predecessor is pure steam.

The technique and methodology of laying out field observations, accounting, analysis and processing of experimental data were carried out according to the generally accepted methodology of B.A. Dospekhov using the Excel computer program (AgCStat) [21, p.245].

The scheme of the experience included the following options:

- 1 – Control;
- 2 – BioSleep BW+Foliar;
- 3 – Orgamica S+Foliar;
- 4 – Organit P+OrganitN+ Biodux +Foliar;
- 5 – Foliar.

The soil of the experimental site is represented by ordinary medium-humus chernozem with a depth of humus horizon of 25-27 cm, and an average humus content of 4,01%. In the arable soil layer of nitrate nitrogen is 14,9 mg, mobile phosphorus is 7,5 mg, and exchangeable potassium is 66,8 mg per 1000 g of soil. Consequently, the nitrogen content is average, phosphorus is low, and potassium is high. According to the mechanical composition, the soil is heavy loamy, the volume weight in the arable horizon is 1,18 g/cm³, and in the meter layer on average is 1,32 g/cm³. The humidity of stable wilting is 12-13%.

Meteorological conditions during the research period were generally typical for the region. Some deviations in moisture and heat reserves corresponded to the definition of a sharply continental climate. Spring in the years of research was dry. According to data from the Chaglinsky meteorological post, 311 mm of atmospheric precipitation fell in the study year of 2022, which is almost equal to the long-term average norm of 315,5 mm (Table 1).

Table 1 – Comparative characteristics of meteorological conditions in alfalfa crops, meteorological post of Shagalaly village (2022-2023)

Years	Month											
	IX	X	XI	XII	I	II	III	IV	V	VI	VII	VIII
Air temperature, °C												
2021-2022	+9,9	+4,3	-6,6	-9,5	-12,4	-9,2	-9,0	+8,5	+13,6	+17,7	+19,9	+16,7
2022-2023	+13	+3,8	-8,3	-15	-12,7	12,6	-3,1	+4,1	+12,7	+18,7	+22,8	+18
Avg. (multi-year average)	+10,8	+3,7	-5,3	-12,7	-16,2	-14,1	-5,9	+4,7	+12,3	+16,8	+18,0	+17,0
Precipitation, mm												
2021-2022	14,2	13,6	18	4,8	11,6	18,5	4,7	5,5	15,7	49,6	77,0	44,1
2022-2023	6,6	37,6	15,8	10,5	12,0	14,0	11,3	1,2	15,8	30,8	20,5	35,5
Avg. (multi-year average)	25,6	25	16,3	12,3	12,4	13,7	16,4	18,8	33,0	43	56,5	37,1

Over the years of research, the distribution of precipitation by month was extremely uneven. The highest amount of precipitation in 2022 was observed in July – 77,0 mm and in June – 49,6 mm, while the lowest figures were in September – 6,6 mm and April – 5,5 mm, which defines the year as dry with an HTC (Hydrothermal Coefficient) of 0,73. The precipitation in June and July positively influenced the formation of the leaf-stem mass of alfalfa plants, while the August precipitation of 44,1 mm positively affected the pod setting of the plants. In the weather conditions of 2023, 300,2 mm of atmospheric precipitation fell, which corresponds to the multi-year average norm of 305,7 mm.

From December to March, 44,5 mm of precipitation fell, compared to the multi-year average of 54,9 mm for this period. A distinctive feature of the autumn-winter period was the high temperature regime. Spring was cold. In March, the average air temperature was -3,1°C. The atmospheric precipitation that fell in March (15,8 mm) did not significantly influence moisture accumulation due to the deficit of pre-winter and winter precipitation. In 2023, the average monthly air temperature in May was +12,7°C, which corresponds to the

multi-year average of +12,4°C, while precipitation in May was 15,8 mm, below the multi-year average of 34,1 mm.

The air temperature in June and July 2023 was higher than the data of 2022 by 1,0°C and 2,9°C, respectively, which positively influenced the growth and development processes of alfalfa plants. Thus, the main indicators – the amount of precipitation and temperature regime – show that the conditions for the growth and development of alfalfa plants in 2023 were satisfactory.

Foliar treatment of crops during vegetation with biological preparations showed different yields in the studied variants. The highest seed yield results were obtained in the second year of life with the use of biological preparations Orgamica S+Foliar -2,72 c/ha, Foliar -2,63 and Organit P, OrganitN, Biodux +Foliar - 2,40 which exceed the control variant without treatment by 0,37 c/ha 0,28 c/ha and 0,05 c/ha respectively. The BioSleep BW+Foliar variant was lower than the control variant by 0,1 c/ha. In the third year of using alfalfa crops, seed yield decreases by 2 times relative to the second year. (Table 2, Figure 1)

Table 2 – The effect of biological preparations on the yield of alfalfa seeds with foliar treatment for vegetation, c/ha

№	Variants	Seed yield, c/ha		
		2022 year	2023 year	average
1	Control	2,35	1,12	1,73
2	BioSleep BW+Foliar	2,25	1,09	1,67
3	Orgamica S+Foliar	2,72	1,29	2,0
4	Organit P,OrganitN, Biodux +Foliar	2,40	1,25	1,82
5	Foliar	2,63	1,10	1,86
	NSR ₀₅	0,24	0,17	0,21




Figure 1 – The effect of biologics on the yield of alfalfa seeds with foliar treatment for vegetation, c/ha

The weight of 1000 seeds are a sign characterizing the quality of the seed material. Large seeds have a larger supply of nutrients than small ones, and as a result, the laboratory and field germination of seeds increases, which is the key to increasing seed yield.

Studies have found that the weight of 1000 seeds for 2022 in the control variant was 2,01 g, and the highest indicator was 2,04 g in the variant of biological preparations Organit P, OrganitN, Biodux +Foliar, treated according to vegetation (Table 3). The coefficient of variability in the experiment by weight of 1000 seeds averaged 3,23%.

Table 3 – The effect of biological preparations on the sowing qualities of seeds with foliar treatment during vegetation

№	Variants	Weight of 1000 seeds, grams			Economic validity of seeds, %		
		2022 year	2023 year	average	2022 year	2023 year	average
1	Control	2,01	1,90	1,95	83,0	75,5	79,2
2	BioSleep BW+Foliar	1,98	1,90	1,94	85,3	75,0	80,1
3	Organica S+Foliar	2,05	2,0	2,02	88,5	80,0	84,2
4	Organit P, OrganitN, Biodux +Foliar	2,02	1,95	1,98	84,5	79,7	82,0
5	Foliar	2,03	2,0	2,0	87,2	80,7	84,0
	V%	2,54	3,42	3,23	2,85	3,42	2,75
	Sx%	2,1	2,6	2,4	2,41	2,64	2,23

Conclusion. The research results confirm two main biological hypotheses of alfalfa culture when cultivated for seeds with foliar top dressing during vegetation:

- firstly, alfalfa generates the highest seed yield in the second year of life, due to the transition to a more complete formation of generative organs. So, if in the second year of plant life, the seed yield is calculated depending on the experimental options from 2,35 to 2,72 c/ha, then in the third year, the level of seed productivity decreases to 1,12 c/ha. Therefore, alfalfa seeds should be left from crops in the second year of life, or at least the third year of life. Then there is a sharp decrease in seed yield and old-age crops do not form economically suitable seed collections;

- secondly, the use of biological products, as one of the main elements of biologized technology, significantly increases seed productivity. So, if the seed yield at the control was 2,35 c/ha, then the use of these biological products increased the yield level to – 2,72 c / ha, that is, by 10-20%.

REFERENCES:

1. Tokusheva A.S., Nugmanov A. B. Problemy' razvitiya kormoproizvodstva v Kazahstane [Problems of feed production development in Kazakhstan]. Agrarlyk gylymdar seriyasy, 2016, pp. 20-40. (In Kazakh).
2. Velikdan N.T. Sostoyanie i perspektivy' razvitiya kormoproizvodstva Stavropol'skogo kraya [The state and prospects of the development of fodder production in the Stavropol Territory]. Vestnik APK Stavropol'ya, 2013, no. 2 (10), pp. 49-53. (In Russian).
3. Kurbanov S. A. Sohranenie i povy'shenie plodorodiya pochv-osnova uvelicheniya effektivnosti zemledeliya Dagestana [Preservation and improvement of soil fertility is the basis for increasing the efficiency of agriculture in Dagestan]. Zemledelie, 2021, no. 4, pp. 16-20. (In Russian).
4. Liu X., Wang, L. LCA-based assessment of hulunber ecological grassland technology integrationdemonstration. Scientia Agricultura Sinica, 2020, vol. 53(13), pp. 2703-2714. doi:10.3864/j.issn.0578-1752.2020.13.018.
5. Zhuchenko A.A. Zernofurazh Rossii [Grain forage of Russia]. Moscow-Kirov, Dom pechaty-Vyatka, 2009, 384 p. (In Russian).
6. Koshen B.M. Agroekologicheskaya ocenka prirodny'h kormovy'h ugodij Severnogo Kazahstana [Agro-ecological assessment of natural forage lands of Northern Kazakhstan]. BBK 74.58 K 75, 2019, 139 p. (In Kazakh).
7. Ombaev A.M., Alimaev I.I. Kormoproizvodstvo-osnova razvitiya zhivotnovodstva Kazahstana [Feed production—the basis for the development of livestock in Kazakhstan]. Privetstvennoe slovo, 2016, 416 p. (In Kazakh).
8. Kosolapov V. M. Strategiya razvitiya selekcii i semenovodstva kormovy'h kul'tur [Strategy of development of the forage crops selection and seed farming]. Kormi i kormovirobnictvo, 2010, no. 67.- pp.3-7. (in Ukraine).
9. Petrina O.V., Braking M. A. Osobennosti razvitiya kornevoj sistemy' lyucerny' sinej i lyucerny' izmenchivoj v usloviyah lesostepnoj zony' srednego urala [Features of the development of the root system of alfalfa and variable alfalfa in the conditions of the forest-steppe zone of the Middle Urals]. BBK 41/4280, 2022, 113 p. (In Russian).
10. Taghizadeh A., Besharati M. Alfalfa properties and livestock nutrition. Alfalfa and clovers. Properties, medicinal uses and health benefits, 2012, pp. 57-78.
11. Solovyova V.N. Vliyanie agropriemov na zasorennost' posevov lyucerny' v usloviyah stepnoj zony' Predural'ya [The influence of agricultural practices on the weed infestation of alfalfa crops in

- the conditions of the steppe zone of the Urals]. *Zhivotnovodstvo i kormoproizvodstvo*, 2013, vol. 3, no.81, pp.127-131. (In Russian).
12. Gubaiddullin H.G. *Lyucerna na korm i semena* [Alfalfa for feed and seeds]. Moscow, Rossel'hozizdat, 1982, 111 p. (In Russian).
 13. Ormeño-Orrillo E., Dávila D.Z. Optimización del tiempo de esterilización de soportes basados en suelo y compost en la producción de inocuentes para leguminosas. *Revista Peruana de Biología*, 1999, vol. 6, no. 2, pp. 181-184. (In Spanish)
 14. Abd-Alla M.H. Synergistic interaction of Rhizobium leguminosarum bv. viciae and arbuscular mycorrhizal fungi as a plant growth promoting biofertilizers for faba bean (*Vicia faba L.*) in alkaline soil. *Microbiological research*, 2014, vol. 169, no. 1, pp. 49-58.
 15. Kneip C. Nitrogen fixation in eukaryotes—new models for symbiosis. *BMC Evolutionary Biology*, 2007, vol. 7, 55 p.
 16. Stepanov A.F., Hramov S. Ispol'zovanie biopreparatov dlya povy'sheniya azotifikiruyushchey sposobnosti i produktivnosti lyucerny' na seroy lesnoy pochve v Zapadnoj Sibiri [The use of biological products to increase the nitrogen-fixing ability and productivity of alfalfa on gray forest soil in western Siberia]. *Vestnik Omskogo Gosudarstvennogo Agrarnogo Universiteta*, 2022, no. 1 (45), pp. 36-44. (In Russian).
 17. Ivanova E.P. Vliyanie defekata i Sinorhizobium meliloti na urozhajnye i kormovy'e dostoinstva lyucerny' izmenchivoj v usloviyah Primorskogo Kraya [The effect of defecate and Sinorhizobium meliloti on the yield and feed advantages of variable alfalfa in the conditions of the Primorsky territory]. *Vestnik Krasnoyarskogo gosudarstvennogo agrarnogo universiteta*, 2023, no. 3 (192), pp. 65-71. (In Russian).
 18. Shahid K., Srivastava V., Sillanpää, M. Protein recovery as a resource from waste specifically via membrane technology – from waste to wonder. *Environmental Science and Pollution Research*, 2021, 28(8), pp.10262-10282.
 19. IFOAM – Organics International. Available at: <https://www.greatitalianfoodtrade.it/ru/mercati/biologico-crescita-piu-lenta-ma-inesorabile-il-rapporto-2023-di-fibl-e-ifoam> (accessed 26 May 2023).
 20. Pochemu lyucerna-novoe zoloto agrobiznesa. [Why alfalfa is the new gold of agribusiness]. Available at: <https://eldala.kz/specproekty/14442> (accessed 06 October 2023). (In Russian).
 21. Dospekhov B.A. Metodika polevogo opy'ta: uchebnik dlya vuzov [Methodology of field experience: manual for the higher educational institutions]. Moscow, Agropromizdat, 1985, 351 p. (In Russian).

Information about the authors:

Kalin Arman Kabdrakhmanuly* – PhD student, "8D08101–Agronomy" educational program, Sh.Ualikhanov Kokshetau University, Republic of Kazakhstan, 020000, Kokshetau, 35 Kudaiberdiyev Str., apt.60, tel.: +77003005694; e-mail: arman.kalin@mail.ru.

Sagalbekov Ualikhan Malgozhdarovich – Doctor of Agricultural Sciences, Professor of the Department of agriculture and bioresources, Sh.Ualikhanov Kokshetau University, Republic of Kazakhstan, 020000, Kokshetau, 76 Abai Str., tel.: +77712784234; e-mail: sagalbekov52@mail.ru.

Auzhanova Mariya Asylkhanovna – Candidate of Agricultural Sciences, acting Associate Professor of the Department of agriculture and bioresources, Sh.Ualikhanov Kokshetau University, Republic of Kazakhstan, 020000, Kokshetau, 76 Abai Str., tel: +7015425375; e-mail: Mayjanova@shokan.edu.kz.

Ualiyeva Gulmira Toktarovna – PhD student, "8D08101 – Agronomy" educational program, Kokshetau experimental production farm, Republic of Kazakhstan, Akmola region, Zerendi district, Shagalaly village, 020000, Akmola region, Kokshetau, 76 Abai Str., tel.: +77051719597; e-mail: ualiyeva_gt@mail.ru.

Калин Арман Қабдракхманұлы* – "8D08101 – Агрономия" мамандығы бойынша докторантуралыңың белім алушысы, Ш. Уәлиханов атындағы Қекшетау университеті, Қазақстан Республикасы, 020000, Қекшетау қ., Құдайбердиеев көшесі, 35, 60-пәтер: тел.+77003005694; е-mail: arman.kalin@mail.ru.

Сагалбеков Уәлихан Малғожарұлы – ауыл шаруашылығы ғылымдарының докторы, ауыл шаруашылығы және биоресурстар кафедрасының профессоры, Ш. Уәлиханов атындағы Қекшетау университеті, Қазақстан Республикасы, 020000, Қекшетау қ., Абай көшесі 76: тел: +77712784234; е-mail: sagalbekov52@mail.ru.

Аужанова Мария Асылхановна – ауыл шаруашылығы ғылымдарының кандидаты, ауыл шаруашылығы және биоресурстар кафедрасының қауымдастырылған профессорының м. а., Ш. Уәлиханов атындағы Қекшетау университеті, Қазақстан Республикасы, 020000, Қекшетау қ., Абай көшесі 76: тел:+7015425375; e-mail: MAUjanova@shokan.edu.kz.

Үәлиева Гулмира Токтарқызы – "8D08101-Агрономия" мамандығының докторанты, Қекшетау тәжірибелік-өндірістік шаруашылығы, Ақмола облысы, Зеренди ауданы, Шағалалы ауылы,

Қазақстан Республикасы, 020000, Көкшетау қ., Абай көшесі 76: тел.+77051719597; e-mail: ualiyeva_gt@mail.ru.

Калин Арман Кабдрахманұлы* – обучающийся докторанттуры по специальности «8D08101 – Агрономия», Кокшетауский университет имени Ш. Уалиханова, Казахстан, 020000, г. Кокшетау, улица Кудайбердиева, 35, кв.60: тел.+77003005694; e-mail: arman.kalin@mail.ru.

Сагалбеков Уалихан Малгождарович – доктор сельскохозяйственных наук, профессор кафедры «Сельское хозяйство и биоресурсы», Кокшетауский университет имени Ш. Уалиханова, Казахстан, 020000, г. Кокшетау, улица Абая 76:т.ел: +77712784234; e-mail: sagalbekov52@mail.ru.

Аужанова Мария Асылхановна – кандидат сельскохозяйственных наук, и.о. ассоциированного профессора кафедры сельское хозяйство и биоресурсы, Кокшетауский университет имени Ш. Уалиханова, Казахстан, 020000, г. Кокшетау, улица Абая 76, тел: +7015425375; e-mail: MAujanova@shokan.edu.kz.

Уалиева Гульмира Токтаровна – докторант специальности «8D08101 – Агрономия», Кокшетауское опытно-производственное хозяйство, Акмолинская область, Зерендинский район, с. Шагалапы, Казахстан, 020000, г. Кокшетау, улица Абая 76: тел.+77051719597; e-mail: ualiyeva_gt@mail.ru.

МРНТИ 34.27.17

УДК 634.8:632

https://doi.org/10.52269/22266070_2024_2_62

ДОМИНИРУЮЩИЕ ВИДЫ НАСЕКОМЫХ-ВРЕДИТЕЛЕЙ, ПОВРЕЖДАЮЩИЕ ДИКОРАСТУЩИЕ ПОПУЛЯЦИИ ЯБЛОНИ СИВЕРСА (*MALUS SIEVERSII*) В ИЛЕЙСКОМ АЛАТАУ

Танабекова Г.Б. – PhD, старший преподаватель кафедры ЮНЕСКО по устойчивому развитию, Казахский национальный университет им. аль-Фараби, старший научный сотрудник лаборатории Энтомологии, Институт Зоологии КН МНВО РК, г. Алматы, Республика Казахстан.

Ященко Р.В.* – д.б.н., генеральный директор, РГП на ПХВ «Институт Зоологии» КН МНВО РК, Научное общество Тетис, г. Алматы, Республика Казахстан.

Лю Жаожи – PhD, профессор ключевых лабораторий Шаньдуна, Сельскохозяйственный университет Циндао, Циндао, Китай.

Статья посвящена проблемам сохранения диких популяций яблони Сиверса (*Malus sieversii*) в Казахстане, которые имеют глобальное мировое значение как природная генетическая основа для поддержания и производства яблок во всем мире. В связи с этим целью публикации является определение степени повреждение яблони Сиверса тремя основными вредителями (*Yponomeuta malinella* Zell., *Archips rosana* L., *Cacoecia crataegana* Hb.) в Илейском Алатау 2018, 2019 годах. Такжедается карта распространения и влияния яблонной горностаевой моли (*Yponomeuta malinella* Zell.), розанной листовертки (*Archips rosana* L.), боярышниковой листовертки (*Cacoecia crataegana* Hb.) на территории Иле-Алатауского ГНПП. Научная значимость исследований заключается в том, что среди основных современных угроз для существования диких популяций яблони Сиверса в последние 20 лет стал местный комплекс видов насекомых-вредителей. Одним из основных условий защиты растений от вредителей является своевременность проведения различных защитных мероприятий. По этой причине изучение современного видового состава, экологических и биологических особенностей доминантных и потенциальных видов насекомых-вредителей яблони Сиверса является неотложной исследовательской задачей, имеющей большое практическое значение.

Ключевые слова: яблоня Сиверса, насекомые-вредители, Илейский Алатау, *Yponomeuta malinella* Zell., *Archips rosana* L., *Cacoecia crataegana* Hb.

ИЛЕ АЛАТАУЫНДАҒЫ СИВЕРС АЛМА АҒАШЫНЫң (*MALUS SIEVERSII*) ЖАБАЙЫ ӨСЕТИН ПОПУЛЯЦИЯЛАРЫН ЗАҚЫМДАЙТАНЫ ЗИЯНКЕС ЖӘНДІКТЕРДІҢ БАСЫМ ТҮРЛЕРІ

Танабекова Г.Б. – PhD, тұрақты даму бойынша ЮНЕСКО кафедрасының аға оқытушы, әл-Фараби атындағы Қазақ ұлттық университеті, Энтомология зертханасының аға ғылыми қызметкері, Зоология институты ҚР ҒжЖБ РК, Алматы қ., Қазақстан Республикасы.

Ященко Р.В.* – б.ғ.д., бас директор, ҚР ҒжЖБ РК Зоология институты, Тетис ғылыми қоғамы, Алматы қ., Қазақстан Республикасы.

Лю Жаожи – PhD, Шаньдунның неаізгі зертханасының профессоры, Циндао ауылшаруашылық университеті, Циндао, Қытай.

Мақала бүкіл әлемде алма жемісін өндіруде табиғи генетикалық негізі ретінде жаһандық маңызы бар Қазақстандағы Сиверс алма ағашының (*Malus sieversii*) жабайы популяцияларын сақтау мәселелеріне арналған. Осыған байланысты басылымның мақсаты Сиверс алма ағашының уш негізгі зиянкестермен (*Yponomeuta malinella* Zell., *Archips rosana* L., *Cacoecia crataegana* Hb.) Iле Алатауында 2018, 2019 жылдары зақымдану дәрежесін анықтая болып табылады. Сондай-ақ, алма ақкіс күйе көбелегінің (*Yponomeuta malinella* Zell.) раушан жапырақ ширатқыш көбелегінің (*Archips rosana* L.) және долана жапырақ ширатқыш көбелегінің (*Cacoecia crataegana* Hb.) таралуы мен әсерінің картасы Iле-Алатау МҰТП аумағында берілген. Зерттеудің ғылыми маңыздылығы соңғы 20 жылда Сиверс алма ағашының жабайы популяцияларының өмір сүруіне негізгі заманауи қауіптердің қатарында зиянкес жәндіктердің жергілікті кешені болып отыр. Өсімдіктерді зиянкестерден қорғаудың негізгі шарттарының бірі – әртүрлі қорғау іс-шараларын үақтылы өткізу. Осы себепті Сиверс алма ағашының зиянкес жәндіктердің басым және ықтимал түрлерінің қазіргі түрлік құрамын, экологиялық және биологиялық ерекшеліктерін зерттеу үлкен практикалық маңызы бар шұғыл зерттеуді қажет етептің міндет болып табылады.

Түйінди сөздер: Сиверс алма ағашы, зиянкес жәндіктер, Iле Алатауы, *Yponomeuta malinella* Zell., *Archips rosana* L., *Cacoecia crataegana* Hb.

DOMINANT INSECT PEST SPECIES THAT DAMAGE WILD POPULATIONS OF THE SIEVERS APPLE TREE (*MALUS SIEVERSI*) IN THE ILE ALATAU

Tanabekova G.B. – PhD, Senior Lecturer of the UNESCO Department on Sustainable Development, Al-Farabi Kazakh National University, Senior Researcher of the Entomology Laboratory, Institute of Zoology CS MSHE RK, Almaty, Republic of Kazakhstan.

Yaschenko R.V.* – Doctor of Biological Sciences, General Director, Institute of Zoology CS MSHE RK, Tetis Scientific Society, Almaty, Republic of Kazakhstan.

Lu Zhaozhi – PhD, Professor of the Shandong Key Laboratories, Qingdao Agricultural University, Qingdao, China.

The article is devoted to the problems of conservation of wild populations of the Sievers apple tree (*Malus sieversii*) in Kazakhstan, which are of global importance as a natural genetic basis for the maintenance and production of apples worldwide. One of the main threats to the existence of wild populations of the Sievers apple tree in the last 20 years has become insect pests that cause enormous damage to natural populations. In this regard, the purpose of the publication is to determine the degree of damage to the Sievers apple tree by three main pests (*Yponomeuta malinella* Zell., *Archips rosana* L., *Cacoecia crataegana* Hb.) in the Ile Alatau in 2018 and 2019. The authors developed a map of the distribution and influence of the apple ermine moth (*Yponomeuta malinella* Zell.), rose leaf roller (*Archips rosana* L.), hawthorn leaf roller (*Cacoecia crataegana* Hb.) on the territory of the Ile Alatau state national natural park. The scientific significance of the research lies in the fact that among the main modern threats to the existence of wild populations of the Sievers apple tree in the last 20 years has become a local complex of insect pest species. One of the main conditions for protecting plants from pests is the timeliness of various events. For this reason, the study of the current species composition, ecological and biological characteristics of dominant and potential insect pests of the Sievers apple tree is an urgent research task of great practical importance.

Key words: Sievers Apple tree, insect pests, Ile Alatau, *Yponomeuta malinella* Zell., *Archips rosana* L., *Cacoecia crataegana* Hb.

Введение.

В число важнейших условий, необходимых для устойчивого развития входит сохранение и рациональное использование биоразнообразия. Актуальность исследования обусловлена тем, что за последние десятилетия ареал обитания этого вида значительно сократился из-за изъятия земель для государственных и экономических нужд, генетического и экологического загрязнения диких популяций, а также появления новых опасных вредителей на окраинных частях ареала.

Яблоня Сиверса является реликтовым видом известным со времен олигоцена и имеющим горно-среднеазиатский тип ареала. Особая ценность вида состоит в том, что он является хранителем уникальной зародышевой плазмы и родоначальником многих культурных сортов яблони [1, с. 23; 2, с. 426]. Естественные насаждения яблони Сиверса не имеют аналогов в мировом растительном сообществе. Они в настоящее время признаны имеющими глобальное мировое значение как единственная в мире природная генетическая основа поддержания и развития культуры яблони [3, с. 305].

Исходя из контекста устойчивого развития, в сохранении яблони Сиверса важно определить основные угрозы и организовать меры по снижению их влияния на природные популяции яблони. В целом, по всему ареалу произрастания яблони Сиверса общая площадь диких популяций этого вида за последние 100 лет сократилась почти на 70%, а генофонд подвергается деградации. Основными